# Specification

## 1.08339.0000 SABOURAUD-2% dextrose broth for microbiology (According harm. EP/ USP/JP)

| | Specification |
|---|---|
| Appearance (clearness) | clear |
| Appearance (color) | yellowish-brown |
| pH-value (25 °C) | 5.4 - 5.8 |

Typical composition (g/litre): Peptone from meat 5.0; Peptone from casein 5.0; D(+)Glucose 20.0.

| | Specification |
|---|---|
| Growth (Trichophyton mentagrophytes ATCC 18748) | good to very good |
| Growth (Trichophyton rubrum ATCC 28188) | fair to good |
| Growth (Microsporum gallinae ATCC 12108) | fair to good |
| Growth (Trichophyton ajelloi ATCC 28454) | fair to good |
| Growth (Aspergillus brasiliensis (formerly A. niger) ATCC 16404 (WDCM 00053)) | good to very good |
| Growth (Penicillium commune ATCC 10428) | good to very good |

Incubation: 7 days; 28°C

| Growth promotion test in accordance with the harmonised method of EP, USP and JP. | Specification |
|---|---|
| Inoculum on reference medium (Candida albicans ATCC 10231 (WDCM 00054)) | ≤ 100 |
| Growth (Candida albicans ATCC 10231 (WDCM 00054)) | fair to very good |

Incubation: up to 3 days; 30-35°C

The information above is current at this time of publication and is subject to change without notice (except for customers holding a change control agreement with our company). The information/format can be altered at any time and inany way if internal company matters or Standard Regulationsdo say so.

Stefanie Fischer
Responsible laboratory manager quality control

This document has been produced electronically and is valid without a signature.